

DVCS and Exclusive Processes at Hermes

Caroline Riedl



Pan-Pacific Symposium on High Energy Physics 2007
Vancouver, Aug. 1, 2007

- Motivation
- GPDs and DVCS Azimuthal Asymmetries
- DVCS Measurements at Hermes
- The Hermes Recoil Detector
- Summarizing Overview

The Composition of the Nucleon's Spin

$$\frac{1}{2} = \underbrace{\frac{1}{2} \Delta\Sigma + L_q}_{J_q} + \underbrace{\Delta G + L_g}_{J_g}$$

- $\Delta\Sigma = 1/3$ from DIS and SIDIS

Hermes: Phys. Rev. **D75** (2007) 012007


$$\Delta\Sigma = 0.330 \pm 0.011 \text{ (theo)} \pm 0.025 \text{ (exp)} \pm 0.028 \text{ (evol)}$$

- ΔG : first indication from DIS and pp \rightarrow small
- $L_q \rightarrow ? \rightarrow$ **Ji's sum-rule!** \leftarrow **Generalized Parton Distributions**
Ji's sum rule: Ji, PRL **78** (1997) 610

$$J_{q,g} = \frac{1}{2} \lim_{t \rightarrow 0} \int_{-1}^1 dx \times [H_{q,g}(x, \xi, t) + E_{q,g}(x, \xi, t)]$$

- $L_g \rightarrow ?$ might be accessible at higher energies than Hermes via GPDs

GPDs: the clever parameterization of the nucleon




unpolarized	polarized	nucleon helicity
$H(x, \xi, t)$	$\tilde{H}(x, \xi, t)$	conserved
$E(x, \xi, t)$	$\tilde{E}(x, \xi, t)$	flipped

- PDFs: $H^q(x, 0, 0) = q(x)$, $\tilde{H}^q(x, 0, 0) = \Delta q(x)$ forward limit
Form Factors: $\int dx [\text{GPD}] = f(t)$, independent of ξ
 \Rightarrow GPDs: simultaneous description of transverse position (FF) and momentum distribution (PDF): "Nucleon Tomography"
- Sum rule for J of quarks/gluons! Need H and E for $t \rightarrow 0$

Recent theoretical reviews:

PPNP **47** (2001) 401; Phys. Rept. **388** (2003) 41; Phys. Rept. **418** (2005) 1

DVCS: the prime process to access GPDs



- $d\sigma \propto |\mathcal{T}|^2 = |\mathcal{T}_{\text{DVCS}}|^2 + |\mathcal{T}_{\text{BH}}|^2 + \mathcal{I}$
- **Interference:** $\mathcal{I} = \mathcal{T}_{\text{DVCS}}\mathcal{T}_{\text{BH}}^* + \mathcal{T}_{\text{DVCS}}^*\mathcal{T}_{\text{BH}}$
- Hermes kinematics: $|\mathcal{T}_{\text{DVCS}}|^2 < |\mathcal{T}_{\text{BH}}|^2$
- $\mathcal{I} \propto \pm(c_0 + \sum_n [c_n \cos(n\phi) + s_n \sin(n\phi)])$
 - ▶ $c_n =$ Lin. Comb. (UU), (UT), (LL), (LT)
 - ▶ $s_n =$ Lin. Comb. (LU), (UL), (UT), (LT)


(beam state, target state) Un-, Long-, Trans.-pol

- Project out sin- and cos-moments by different:
 - ▶ beam charges
 - ▶ beam helicities
 - ▶ target polarizations (long. and trans.)

\Rightarrow Azimuthal asymmetries


\Rightarrow Linear combinations of GPDs

The Hermes forward spectrometer




The powerful DVCS data pool of Hermes

Hermes (1): 1996-2000, (2): 2002-2005, (3): 2006-2007



- 2 beam charges (1,2,3), 2 beam helicities (1,2,3)
- longitudinally (1) and transversely (2) polarized target
- unpolarized nuclear targets (1,2)
- Recoil Detector (3)

Finding exclusive events



sidis background: 5%, associated: 11%

- Classic technique at Hermes: access to exclusivity via missing mass

$$M_x^2 = (P_e + P_P - (P_{e'} + P_\gamma))^2$$

- 2006/07: recoiling proton detected \Rightarrow all reaction partners measured!

The gallery of DVCS azimuthal asymmetries

1. Beam Charge Asymmetry $A_C(\phi)$

$$d\sigma(e^+, \phi) - d\sigma(e^-, \phi) \propto \text{Re}(F_1 \mathcal{H}) \cos \phi$$

2. Beam Spin Asymmetry $A_{LU}(\phi)$


$$d\sigma(\vec{e}, \phi) - d\sigma(\overleftarrow{e}, \phi) \propto \text{Im}(F_1 \mathcal{H}) \sin \phi$$

3. Longitudinal Target Spin Asymmetry $A_{UL}(\phi)$

$$d\sigma(\overleftarrow{P}, \phi) - d\sigma(\overrightarrow{P}, \phi) \propto \text{Im}(F_1 \tilde{\mathcal{H}}) \sin \phi$$

4. Transverse Target Spin Asymmetry $A_{UT}(\phi, \phi_s)$

$$d\sigma(\phi, \phi_s) - d\sigma(\phi, \phi_s + \pi) \propto \text{Im}(F_2 \mathcal{H} - F_1 \mathcal{E}) \sin(\phi - \phi_s) \cos \phi \\ + \text{Im}(F_2 \tilde{\mathcal{H}} - F_1 \xi \tilde{\mathcal{E}}) \cos(\phi - \phi_s) \sin \phi$$




DVCS-BH interference term \mathcal{I} induces azimuthal asymmetries \Rightarrow GPDs

F_1, F_2 : PAULI, DIRAC Form Factors; $\mathcal{H}, \tilde{\mathcal{H}}, \mathcal{E}, \tilde{\mathcal{E}}$: COMPTON Form Factors
(convolutions of hard scattering amplitude and corresponding twist-2 GPD)

1. Beam Charge Asymmetry (BCA) versus ϕ

$$A_C(\phi) = \frac{d\sigma(e^+, \phi) - d\sigma(e^-, \phi)}{d\sigma(e^+, \phi) + d\sigma(e^-, \phi)} \propto \text{Re}(F_1 \mathcal{H}) \cos \phi$$




$$P_1 = -0.011 \pm 0.019$$

$$P_2 = 0.060 \pm 0.027$$

$$P_3 = 0.016 \pm 0.026$$

$$P_4 = 0.034 \pm 0.027$$

BCA $\cos\phi$ amplitude





- Model (proton data only):
PRD60 (1999) 094017 and
PPNP47 (2001) 401 \Rightarrow
Regge-inspired with D-term
disfavored
- Contributions for $ed \rightarrow eX\gamma$:
coherent ($X=d$): 20%
incoherent ($X=pn$): 60%
associated ($X=\Delta$): 15%

Hermes publication: Phys. Rev. **D75** (2007) 011103(R)

Factor of ≈ 20 more e^- and factor of ≈ 7 more e^+ data on tape!

2. Beam Spin Asymmetry (BSA)

$$A_{LU}(\phi) = \frac{1}{\langle |P_B| \rangle} \cdot \frac{d\sigma(\vec{e}, \phi) - d\sigma(\overleftarrow{e}, \phi)}{d\sigma(\vec{e}, \phi) + d\sigma(\overleftarrow{e}, \phi)} \propto \text{Im}(F_1 \mathcal{H}) \sin \phi$$





Hermes publication: PRL **87** (2001) 182001

Factor of ≈ 9 more data on tape

3. Longitudinal Target Spin Asymmetry (LTSA)

$$A_{UL}^{\sin\phi}(\phi) = \frac{1}{\langle |P_T| \rangle} \cdot \frac{d\sigma(\overleftarrow{P}, \phi) - d\sigma(\overrightarrow{P}, \phi)}{d\sigma(\overleftarrow{P}, \phi) + d\sigma(\overrightarrow{P}, \phi)} \propto \text{Im}(F_1 \tilde{\mathcal{H}}) \sin\phi$$




GPD model: see reference @ BCA

Plots: Full statistics

4. Transverse Target Spin Asymmetry (TTSA)

$$A_{UT}(\phi, \phi_s) = \frac{1}{\langle |P_T| \rangle} \cdot \frac{d\sigma(\phi, \phi_s) - d\sigma(\phi, \phi_s + \pi)}{d\sigma(\phi, \phi_s) + d\sigma(\phi, \phi_s + \pi)} \propto$$



sensitive to J_U :

$$\text{Im}(F_2 \mathcal{H} - F_1 \mathcal{E}) \cdot \sin(\phi - \phi_s) \cos \phi +$$

NOT sensitive to J_U :


$$\text{Im}(F_2 \tilde{\mathcal{H}} - F_1 \xi \tilde{\mathcal{E}}) \cdot \cos(\phi - \phi_s) \sin \phi$$

Sensitivity on J_U : hep-ph/0506264, assuming $J_d = 0$

Factor of 1 more data

First (Model-Dependent) Constraint on $J_u + k \cdot J_d$

$$\chi^2(J_u, J_d) = \frac{\left(A_{\text{UT}}^{\sin(\phi-\phi_s) \cos \phi} \Big|_{\text{exp}} - A_{\text{UT}}^{\sin(\phi-\phi_s) \cos \phi} \Big|_{\text{VGG}(J_u, J_d)} \right)^2}{\delta A_{\text{stat}}^2 + \delta A_{\text{sys}}^2}$$



- J_u and J_d free params in GPD model (VGG)

- 1-sigma constraint on

J_u vs. J_d :

$$\chi^2(J_u, J_d) \leq \chi_{\text{min}}^2 + 1$$


(brown band)

2b. DVCS on nuclear targets

- How does the nuclear environment modify parton-parton correlations?
- **Hermes nuclear targets:** ^2H , ^4He , ^{14}N , ^{20}Ne , $^{82-86}\text{Kr}$, $^{129-134}\text{Xe}$

2b. DVCS on nuclear targets

- How does the nuclear environment modify parton-parton correlations?
- **Hermes nuclear targets:** ^2H , ^4He , ^{14}N , ^{20}Ne , $^{82-86}\text{Kr}$, $^{129-134}\text{Xe}$
- Nuclear BSA: **clear $\sin\phi$ amplitude in the exclusive region**




Integrated kinematics:

Neon: $-0.22 \pm 0.03 \pm 0.03$

proton: $-0.18 \pm 0.03 \pm 0.03$

2b. DVCS on nuclear targets

- How does the nuclear environment modify parton-parton correlations?
- **Hermes nuclear targets:** ^2H , ^4He , ^{14}N , ^{20}Ne , $^{82-86}\text{Kr}$, $^{129-134}\text{Xe}$



Nuclear-to-hydrogen ratio
of BSA $\sin \phi$ amplitudes:

$$(A/H) = 1.58 \pm 0.26$$

(coherent)


incoherent:

(A/H) consistent with 1
✓ GPD model prediction


PRC68 (2003) 015204

Factor of $\approx 2/1$ more data on tape for Xenon/Krypton

Recoil Detector installation: December 2005





The Hermes Recoil Detector



- Superconducting Solenoid (1 Tesla)
- Photon Detector
 - ▶ 3 layers of Tungsten/Scintillator
 - ▶ π^0 **background supression**
- Scintillating Fiber Tracker
 - ▶ 2 Barrels
 - ▶ Each 2 parallel- & 2 stereo-layers
 - ▶ Stereo angle: 10°
 - ▶ **Momentum reconstruction & PID**
- Silicon Strip Detector
 - ▶ 2 Layers
 - ▶ 16 double-sided sensors
 - ▶ (10cm \times 10cm) active area
 - ▶ Inside accelerator vacuum
 - ▶ **Momentum reconstruction & PID**
- Target Cell with unpol. H_2 or D_2

The Recoil Road to genuine exclusivity at Hermes



Silicon & Fiber Tracker:

$$p_p \in [135, 1200] \text{ MeV}/c$$

p/π PID for $p < 700 \text{ MeV}/c$

Photon Detector:

p/π PID for $p > 650 \text{ MeV}/c$


- Recoiling protons
 - Enhance **signal fraction**
 - Improve t -resolution
- Background pions and protons
- Photons from $\pi^0 \rightarrow \gamma\gamma$

⇒ Reduce **background contributions:**


– **sidis:** 5% $\searrow \ll 1\%$

– **associated production:** 11% $\searrow 1\%$

Recoil: First physics signatures




Recoil: First physics signatures





Momentum (p) reconstruction:

- 1 Low momentum protons:
 $\Rightarrow p$ by $\Sigma(\text{energy deposits})$
- 2 Higher momentum protons:
 $\Rightarrow p$ by Bethe-Bloch (dE/dx)
- 3 High momentum particles:
 $\Rightarrow p$ by bending in B-field,
 \Rightarrow tracks formed by
spacepoints in (up to) 2
subdetectors

Recoil: First physics signatures





⇒ p/π PID:



Recoil: Proof of Principle

Response of Recoil Silicon Detector for “traditional DVCS candidates”
(events with 1 lepton and 1 photon in front spectrometer):




The Recoil Adventure goes on!

- Hera stopped its operation 4 weeks ago, Hermes took last data
- Collected statistics (*preliminary*) with operational recoil detector:
 - ▶ **Electron beam** 2006 (only Fiber Tracker operational):
 H_2 : 5k DVCS (3 Mio DIS), D_2 : 1k DVCS (0.8 Mio DIS)
 - ▶ **Positron beam** 2006/07 (all subdetectors fully operational):
 H_2 : 42k DVCS (28 Mio DIS), D_2 : 10k DVCS (7 Mio DIS)
- Analysis of BCA and BSA with recoil data
- Detector understanding in progress
 - ▶ Calibration
 - ▶ Alignment
 - ▶ Noise
 - ▶ Tracking
 - ▶ ...

⇒ Watch out for first results!

Summary: Hermes and Exclusive Processes



unpolarized	polarized	
H : BCA, BSA, TTSA E : TTSA	\tilde{H} : LTSA (TTSA) \tilde{E} : (TTSA)	
$J^P = 1^-$ mesons	x-section $J^P = 0^-$ mesons	see talk J. Dreschler
photon: $J^P = 1^-$ (DVCS)		

- GPD models agree in general with measurements
- First model-dependent extraction of $J_u + k \cdot J_d$ possible
- Most published DVCS results await a significant statistics upgrade: BCA (factor 20 / 7 more), BSA (factor 9), TTSA (factor 1)
- Recoil-data is being prepared for physics analysis
 \Rightarrow exploit direct exclusivity: no mass is missing anymore!
- Once background contribution is measured: refined analysis of pre-recoil DVCS and DVMP data